arXiv:1610.07356v1 [math.GT] 24 Oct 2016

NESTED OPEN BOOKS AND THE BINDING SUM

SEBASTIAN DURST AND MIRKO KLUKAS

Abstract. We introduce the notion of a *nested open book*, a submanifold equipped with an open book structure compatible with an ambient open book, and describe in detail the special case of a *push-off* of the binding of an open book. This enables us to explicitly describe a natural open book decomposition of a fibre connected sum of two open books along their bindings, provided they are diffeomorphic and admit an open book structure themselves. Furthermore, we apply the results to contact open books, showing that the natural open book structure of a contact fibre connected sum of two adapted open books along their contactomorphic bindings is again adapted to the resulting contact structure.

Contents

1. Introduction 1
1.1. Organisation of the paper 3
2. Preliminaries 3
2.1. Open book decompositions 3
2.2. The fibre connected sum and the binding sum 5
2.3. Generalised 1-handles 7
3. Nested open books 7
3.1. Fibre sums along nested open books 8
4. The push-off 10
4.1. Framings 11
4.2. The push-off as an abstract nested open book 14
5. Proof of Theorem 1.1 17
5.1. Proof of the 1st part: Existence and identifying the monodromy 18
5.2. Proof of the 2nd part: Handle interpretation of the page 18
6. Application to contact topology 20
6.1. Proof of Theorem 1.3 23
6.2. Recipe for an abstract description 24
References 26

1. Introduction

In 1923 Alexander [1] proved that every closed oriented 3-manifold admits a so-called *open book decomposition*. In fact, combining the work of Winkelnkemper, Lawson, and Quinn from the 1970s, this statement remains true for odd-dimensional manifolds in general (see [18], [12], and [14] respectively). The existence problem

Date: October 25, 2016.

in even dimensions is also solved in these works but is more involved. Arguably the most prominent appearance of open books in recent history is their surprisingly deep connection with *contact structures*. As was observed by Giroux [7] in 2002, contact structures in dimension 3 are of purely topological nature: he established a one-to-one correspondence between isotopy classes of contact structures and open book decompositions up to positive stabilisation. This correlation remains partly intact in higher dimensions. According to Giroux and Mohsen [9] any contact structure on a closed manifold of dimension at least 3 admits a *compatible* open book decomposition.

In the present paper we investigate how the *fibre connected sum* performed along diffeomorphic binding components of an open book, henceforth called *binding sum*, affects an underlying open book structure. We discuss both regular fibre sums and fibre sums of contact manifolds. While, at first glance, the binding sum destroys the open book structure, we show that the resulting manifold again admits an open book decomposition, provided the binding components admit open book structures themselves. Furthermore, we describe an explicit open book structure in terms of the original open book, thus generalising a previous result of the second author [10] to higher dimensions. More concretely, we give a constructive proof of the following result.

Theorem 1.1. *Let M be a (not necessarily connected) smooth manifold with open book decomposition (Σ, ϕ) whose binding B contains two diffeomorphic components B_1, B_2 with diffeomorphic open book decompositions (Σ', ϕ'). Then the fibre connected sum of M performed along B_1 and B_2 with respect to the page framings admits an open book decomposition. The new page can be can be obtained from Σ by two consecutive generalised* 1*-handle attachments whose type depends on the pages Σ' of the binding components*[1]*. The new binding is given by a fibre connected sum of B_1 and B_2 along their respective bindings $B' = \partial\Sigma$. Away from the handle attachments the monodromy remains unchanged, and over the remaining part it restricts to $\psi \circ \mathcal{D}$, where ψ is a* Chinese burn *along B_i (see Definition 4.4) and $\mathcal{D}$ the* twist map *(see Definition 4.7).*

Remark 1.2. The statement of Theorem 1.1 still holds if the two binding components admit similar fibrations over the circle (i.e. , informally, if their respective nested bindings are empty). In particular, the statement holds in the 3-dimensional case, where the binding components are circles.

Note that, according to the above mentioned work of Winkelnkemper, Lawson, and Quinn, the condition on the binding components to admit open books themselves is not a restriction in odd dimensions. For the same reasons the sheer existence of an open book of the fibre sum in odd dimensions is already answered as well. According to Giroux and Mohsen the same is true for the case of adapted open books and contact structures. However, constructions of particular instances of those open books and their relationship to the original open book decomposition have not been discussed in the literature yet. Besides, very few constructions of open books supporting contact structures in higher dimensions are known. Along these lines we prove that our construction can also be adapted to an underlying contact structure and again produces a compatible open book.

[1]Let Σ' be a page of the open book for B, then a generalized 1-handle in the sense of [11] is of the form $H_{\Sigma'} = D^1 \times (\Sigma' \times D^1)$.

Theorem 1.3. *Let (M,ξ) be a (not necessarily connected) contact manifold carried by an open book (B,π), and assume that the binding B equipped with the induced contact structure contains two contactomorphic components B_0 and B_1. Then their push-offs B_i^+ are contact submanifolds contact isotopic to the respective binding component B_i. Furthermore, the natural contact structure on the contact fibre connected sum along the B_i^+ is supported by the resulting open book.*

1.1. **Organisation of the paper.** After recalling some basic facts about open book decompositions and fibre sums we introduce the notion of a *nested open book* in Section 3, discussing fibre connected sums in this context. We will then proceed to treat in detail the special case of the *binding sum*. Section 4 describes an appropriately framed *push-off* of the binding of an open book, realizing it as a nested open book, and hence presenting a better candidate to perform the fibre sum. The main result is stated in detail and proved in Section 5. We conclude the topological part of the paper by interpreting the binding connected sum as two consecutive generalized 1-handle attachments in Section 5.2 before turning our attention to the adaptation of the constructions to contact topology in Section 6.

Remark 1.4. All manifolds are understood to be smooth.

2. Preliminaries

2.1. **Open book decompositions.** An **open book decomposition** of an n-dimensional manifold M is a pair (B,π), where B is a co-dimension 2 submanifold in M, called the **binding** of the open book, and $\pi\colon M\setminus B\to S^1$ is a (smooth, locally trivial) fibration such that each fibre $\pi^{-1}(\theta)$, $\theta\in S^1$, corresponds to the interior of a compact hypersurface $\Sigma_\theta\subset M$ with $\partial\Sigma_\theta = B$. The hypersurfaces Σ_θ, $\theta\in S^1$, are called the **pages** of the open book.

In some cases we are not interested in the exact position of the binding or the pages of an open book decomposition inside the ambient space. Therefore, given an open book decomposition (B,π) of an n-manifold M, we could ask for the relevant data to remodel the ambient space M and its underlying open book structure (B,π), say up to diffeomorphism. This leads us to the following notion.

An **abstract open book** is a pair (Σ,ϕ), where Σ is a compact hypersurface with non-empty boundary $\partial\Sigma$, called the **page**, and $\phi\colon\Sigma\to\Sigma$ is a diffeomorphism equal to the identity near $\partial\Sigma$, called the **monodromy** of the open book. Let $\Sigma(\phi)$ denote the mapping torus of ϕ, that is, the quotient space obtained from $\Sigma\times[0,2\pi]$ by identifying $(x,2\pi)$ with $(\phi(x),0)$ for each $x\in\Sigma$. Then the pair (Σ,ϕ) determines a closed manifold $M_{(\Sigma,\phi)}$ defined by

$$M_{(\Sigma,\phi)} := \Sigma(\phi)\cup_{\mathrm{id}}(\partial\Sigma\times D^2), \tag{2.1}$$

where we identify $\partial\Sigma(\phi)=\partial\Sigma\times S^1$ with $\partial(\partial\Sigma\times D^2)$ using the identity map. Let $B\subset M_{(\Sigma,\phi)}$ denote the embedded submanifold $\partial\Sigma\times\{0\}$. Then we can define a fibration $\pi\colon M_{(\Sigma,\phi)}\setminus B\to S^1$ by

$$\left.\begin{matrix}[x,\theta]\\ [x',re^{i\theta}]\end{matrix}\right\}\mapsto[\theta],$$

where we understand $M_{(\Sigma,\phi)}\setminus B$ as decomposed as in (2.1) and $[x,\theta]\in\Sigma(\phi)$ or $[x',re^{i\theta}]\in\partial\Sigma\times D^2\subset\partial\Sigma\times\mathbb{C}$. Clearly, (B,π) defines an open book decomposition of $M_{(\Sigma,\phi)}$.



FIGURE 1. Schematic picture of an open book. A single page of a nested open book and its nested binding is indicated.

On the other hand, an open book decomposition (B, π) of some n-manifold M defines an abstract open book as follows: identify a neighbourhood of B with $B \times D^2$ such that $B = B \times \{0\}$ and such that the fibration on this neighbourhood is given by the angular coordinate, θ say, on the D^2-factor. We can define a 1-form α on the complement $M \setminus (B \times D^2)$ by pulling back $d\theta$ under the fibration π, where this time we understand θ as the coordinate on the target space of π. The vector field $\partial\theta$ on $\partial\big(M \setminus (B \times D^2)\big)$ extends to a nowhere vanishing vector field X which we normalise by demanding it to satisfy $\alpha(X) = 1$. Let ϕ denote the time-2π map of the flow of X. Then the pair (Σ, ϕ), with $\Sigma = \overline{\pi^{-1}(0)}$, defines an abstract open book such that $M_{(\Sigma,\phi)}$ is diffeomorphic to M.

Remark 2.1 (Existence of open book decompositions)**.** In 1923 Alexander [1] proved that every closed oriented 3-manifold admits an open book decomposition. Winkelnkemper [18] showed that a closed oriented simply-connected manifold of dimension at least six can be given the structure of an open book if and only if its signature vanishes. In particular, that is the case in odd dimensions by definition. For odd-dimensional manifolds the hypothesis of simply-connectedness can be dropped, as was shown by Lawson [12]. Quinn [14] extended this to 5-manifolds and also discussed the even-dimensional case, where the obstruction is more involved than in the simply-connected case.

Example 2.2. A 2-dimensional open book has to have closed intervals as pages and thus trivial monodromy. Hence, the only surface admitting an open book decomposition is the sphere and the decomposition is unique up to isotopy.

Example 2.3. The 3-dimensional sphere S^3 has an open book decomposition with page a disc and trivial monodromy. This can be seen by realizing S^3 as the one-point compactification of $\mathbb{R}^3$ and extending the binding and pages in Figure 1 along the z-axis. To be more precise, choosing polar coordinates (r, θ) on the xy-plane we define $B := \{x = y = 0\} \cup \{\infty\}$ and $\pi\colon\ S^3 \setminus B \to S^1$ by sending (r, θ, z) to θ, where we identify $S^3 \equiv \mathbb{R}^3 \cup \{\infty\}$.

Example 2.4. Consider S^3 as the unit sphere in $\mathbb{C}^2$ and denote the positive Hopf link by $B := \{(z_1, z_2) \in S^3 : z_1 z_2 = 0\}$. Then

$$\pi\colon\ S^3 \setminus B \to S^1 \subset \mathbb{C},\ (z_1, z_2) \mapsto \frac{z_1 z_2}{|z_1 z_2|}$$

defines an open book decomposition with page an annulus and monodromy a positive Dehn-twist along the core of the annulus.

2.1.1. *Open books and handlebodies.* An open book decomposition (B, π) yields a decomposition of the underlying manifold M into two diffeomorphic solid handlebodies, where each corresponds to "one half of the open book":

$$M = \overline{\pi^{-1}([0, \pi])} \cup \overline{\pi^{-1}([\pi, 2\pi])}.$$

In particular, the handlebodies in this splitting are the quotient of the product of the page Σ of the open book with the interval, where each $\{p\} \times [0, \pi]$ for $p \in \partial\Sigma$ is smashed to a point. Conversely, certain decompositions of a manifold as a double of a handlebody $M = H_1 \cup H_2$ give rise to an open book structure: Assume that ∂H_i is given the structure of a cell complex and there exists some embedded subcomplex $K \subset \partial H_i$ such that the inclusions into H_i are homotopy equivalences and such that H_i is diffeomorphic to the product of the interval I with a regular neighbourhood V of K in ∂H_i. Then M has an open book decomposition with page V.

Example 2.5. The genus 1 Heegaard splitting of the 3-sphere induces the open book decomposition described in Example 2.4: We need K on the torus such that the inclusions into both solid tori in the Heegaard decomposition are homotopy equivalences. This forces K to be a $(1, 1)$-torus knot. A regular neighbourhood of K on the Heegaard torus is an annulus. The induced monodromy is a Dehn twist.

Remark 2.6. Note that it is not always possible to write a handlebody as a product of this special form. Consider the connected sum of two copies of $\mathbb{C}P^4$. This has a handle decomposition consisting of a single 0- and 8-handle and two handles of every even index in between. Thus, we can write $\mathbb{C}P^4 \# \mathbb{C}P^4$ as the union of two diffeomorphic solid handlebodies, each consisting of a single 0- and 4-handle and two 2-handles. However, $\mathbb{C}P^4 \# \mathbb{C}P^4$ has non-vanishing signature, so it does not admit an open book decomposition by the results mentioned in Remark 2.1.

Nice surveys on open books and their applications are Winkelnkemper's appendix to Ranicki's book [15] and Giroux [8]. More detailed material, in particular on the relation of open books and contact structures can be found in [5, 6, 16].

2.2. **The fibre connected sum and the binding sum.** We will briefly introduce the fibre connected sum, which is a method to construct manifolds using embedded submanifolds; for details see [6, Section 7.4]. Let M' and M be closed oriented manifolds and let j_0 and j_1 be embeddings of M' into M with disjoint images. Assume that there exists a bundle isomorphism Ψ of the corresponding normal

bundles N_0 and N_1 over $j_1 \circ j_0^{-1}|_{j_0(M')}$ that reverses the fibre orientation. Picking a bundle metric on N_0 and choosing the induced metric on N_1 turns Ψ into a bundle isometry. We furthermore identify open disjoint neighbourhoods of the $j_i(M')$ with the normal bundles N_i.

The **fibre connected sum** is the quotient manifold

$$\#_\Psi M := \Big(M \setminus \big(j_0(M') \cup j_1(M')\big)\Big)/\sim,$$

where $v \in N_0$ with $0 < \|v\| < \epsilon$ is identified with $\frac{\sqrt{\epsilon^2 - \|v\|^2}}{\|v\|}\Psi(v)$. A useful interpretation of the fibre sum is the following: suppose we identify the the boundaries of the embedded normal bundles N_0 and N_1 with their induced sphere bundles. Then the fibre connected sum is diffeomorphic to the quotient

$$\Big(M \setminus \big(N_0 \cup N_1\big)\Big)/\sim,$$

where we identify $p \in \partial N_0$ with $\Psi(p) \in \partial N_1$. Observe that (in both cases) the identification is orientation preserving. If M is disconnected and j_i maps M' into M_i with $M = M_0 \sqcup M_1$, we also write $M_0 \#_\Psi M_1$ for $\#_\Psi M$. In the case when M' is just a point the fibre connected sum coincides with the ordinary connected sum. Note also that there is a cobordism from M to $\#_\Psi M$, i.e. the fibre connected sum can be obtained by a sequence of surgeries. Furthermore, it is worth noting that the construction can be adapted to work in the symplectic and contact setting if the dimensions of M and M' differ by two, see Section 7.2 in [13] and Theorem 7.4.3 in [6] for details.

Let M be a (not necessarily connected) smooth n-dimensional manifold with open book decomposition (Σ, ϕ) whose binding B contains two diffeomorphic components B_1, B_2 with diffeomorphic open book decompositions (Σ', ϕ'). Their normal bundles νB_0 and νB_1 admit trivializations induced by the pages of the open book decomposition of M. Let Ψ denote the fibre orientation reversing diffeomorphism of $B \times D^2 \subset B \times \mathbb{C}$ sending (b, z) to $(b, \bar{z})$. Hence, we can perform the fibre connected sum along B_1 and B_2 with respect to the above trivializations of the normal bundles and the map induced by Ψ and denote the result by

$$\#_{B_0, B_1} M.$$

We call it the **binding sum** of M along B_1 and B_2.

An interesting case occurs when M is disconnected and decomposes into two manifolds M_0 and M_1 with open books admitting diffeomorphic bindings, denoted by B say. In this case we denote the binding sum along the receptive copies of B in M_0 and M_1 by

$$M_0 \#_B M_1.$$

Note that $M_0 \#_B M_1$ admits the structure of a fibration over the circle with fibre given by

$$(-\Sigma_0) \cup_B \Sigma_1,$$

where Σ_0 and Σ_1 are the pages of the open book for M_0 and M_1 respectively. In the contact setting each fibre defines a *convex* hypersurface, i.e. there exists a contact vector field on $(M_0, \xi_0) \#_B (M_1, \xi_1)$ which is transverse to the fibres. Furthermore for each fibre $(-\Sigma_0) \cup_B \Sigma_1$ the contact vector field is tangent to the contact structure exactly over B.

This paper is concerned with the question whether the binding sum admits again the structure of an open book and how it is related to the original open books.

2.3. **Generalised 1-handles.** In this section we will briefly introduce generalised 1-handles in the sense of [11] and their attachment. Recall that an ordinary n-dimensional 1-handle is of the form $D^1 \times D^{n-1}$, and note that we may understand the $(n-1)$-dimensional disc as a thickened disc of dimension $n-2$. We are now simply going to replace this disc by an appropriate manifold with boundary.

Let Σ be an $(n-2)$-dimensional manifold with boundary. An **n-dimensional generalised 1-handle** is given by

$$H_\Sigma := D^1 \times (\Sigma \times D^1).$$

Its boundary decomposes into two parts, namely

$$\partial_- H_\Sigma = S^0 \times (\Sigma \times D^1) \quad \text{and} \quad \partial_+ H_\Sigma = D^1 \times \partial(\Sigma \times D^1).$$

In analogy to a regular 1-handle we define the **core** of H_Σ to be $D^1 \times (\Sigma \times \{0\})$ and its **co-core** to be $\{0\} \times (\Sigma \times D^1)$. We **attach** a generalised 1-handle H_Σ to an n-dimensional manifold M with boundary ∂M via an embedding $f\colon \partial_- H_k \to \partial M$ to obtain a manifold $M' = M \cup_f H_\Sigma$. Corners are understood to be smoothened (see Definition und Notiz (13.12) in [2]).

Attaching generalised handles with isotopic maps f_0 and f_1 results in diffeomorphic manifolds $M_0 = M \cup_{f_0} H_\Sigma$ and $M_1 = M \cup_{f_1} H_\Sigma$. An isotopy of the attaching maps defines a time-dependent vector field on its image that can be extended to the whole manifold. The corresponding time-1 map is the desired diffeomorphism. Observe that to define a generalised handle attachment it is sufficient to specify the image of the boundary $S^0 \times \Sigma \times \{0\}$ of the core of the handle under the attaching map.

It is worth noting that, in case Σ can be endowed with an appropriate exact symplectic form, the above construction can be adapted to the contact setting, and naturally extends the symplectic handle constructions due to Eliashberg [4] and Weinstein [17].

3. Nested open books

In this section we turn our attention to a special class of submanifolds and introduce the notion of a *nested open book*, i.e. a submanifold carrying an open book structure compatible with the open book structure of the ambient manifold. We also discuss fibre connected sums in this context.

Let M be an n-dimensional manifold supported by an open book decomposition (B, π). Let $M' \subset M$ be a k-dimensional submanifold on its part supported by an open book decomposition (B', π') such that

$$\pi|_{M' \setminus B'} = \pi'.$$

Note that B' necessarily defines a $(k-2)$-dimensional submanifold in B. We will always assume that M' intersects the binding B transversely. We refer to M', as well as to (B', π'), as a **nested open book** of (B, π).

Remark 3.1. A nested open book can be compared to the images in a *flip-book*: every page of the ambient open book contains a nested page, and these nested pages "move" like the drawings in a flip-book when flipping through its pages.

Let (Σ, ϕ) be an abstract open book and $\Sigma' \subset \Sigma$ a properly embedded submanifold with boundary $\partial\Sigma' \subset \partial\Sigma$. We call $(\Sigma', \phi|_{\Sigma'})$ an **abstract nested open book** if Σ' is invariant under the monodromy ϕ. The equivalence of the two definitions follows analogously to the equivalence of abstract and non-abstract open books. If not indicated otherwise, we will assume the normal bundle of any nested open book used in the present paper to be trivial.

Example 3.2. Consider a k-disc $D^k \subset D^n$ inside an n-disc D^n coming from the natural inclusion $\mathbb{R}^k \subset \mathbb{R}^n$. This realises $S^{k+1} \cong (D^k, \mathrm{id})$ as a nested open book of $S^{n+1} \cong (D^n, \mathrm{id})$. The case $k = 1$ and $n = 2$ is depicted in Figure 1. For $k = n - 2$, the nested S^{n-1} is a *push-off*, as will be defined in Section 4, of the binding of (D^n, id); cf. also Example 5.1, where we discuss the binding sum of two copies of (D^3, id).

3.1. **Fibre sums along nested open books.** For the remainder of the section, assume the co-dimension of the nested open books to be two. The question whether the resulting manifold of a fibre connected sum operation of two open books carries an open book structure can be answered positively in the case when the sum is performed along nested open books. The page of the resulting open book is a fibre connected sum of the pages (along the page of the nested book), the binding is a fibre connected sum of the bindings (along the nested binding), and the monodromies glue together. In Section 4 we will consider a special case in detail and discuss an explicit open book structure of the sum, namely, the case of the nested open book being a push-off of the binding. Not only is it possible to describe the monodromy in terms of the old monodromies, but we also obtain a description of the page via generalised handle attachments. This description of the new page is not always possible in the case of arbitrary nested books, since the nested page can be entangled, i.e. cannot be isotoped to the boundary, inside the ambient page. Let us first study the situation of fibre sums along nested open books in general.

Let M' be an $(n-2)$-dimensional manifold supported by an open book (π', B'), and let $j_0, j_1 \colon M' \hookrightarrow M$ be two disjoint embeddings defining nested open books of M such that their images admit isomorphic normal bundles. We denote by $M'_i := j_i(M')$ the embedded copies of the nested open book M' and by $B'_i := j_i(B')$ their respective bindings. Finally let $\pi'_i := \pi' \circ j_i^{-1}$ denote the induced open book structure on $M'_i \setminus B'_i$ and let $(\Sigma'_i)_\theta$ denote their pages.

Given an orientation reversing bundle isomorphism Ψ of the normal bundles $\nu M'_i$, we can perform the fibre connected sum $\#_\Psi M$. We only have to ensure that the fibres of the normal bundles of M'_0 and M'_1 lie within the pages of (π, B). In particular, we require the fibres over the nested bindings to lie within the binding of M. Moreover, we require the isomorphism Ψ of the normal bundle to respect the open book structure of M (in particular this implies that it is compatible with the nested open book structures of M'_0 and M'_1), i.e. we require Ψ to satisfy $\pi \circ \Psi = \pi$. Now, an open book structure of $\#_\Psi M$ is given as follows.

Lemma 3.3. *The original fibration $\pi \colon M \setminus B \to S^1$ descends to a fibration*

$$\Pi \colon \#_\Psi M \setminus \#_{\Psi|_{\nu B'_0}} B \to S^1.$$

In particular, the new binding is given by the fibre connected sum $\#_{\Psi|_{\nu B'_0}} B$ of the binding along the nested bindings (with respect to the isomorphism of $\nu B'_i \subset TB$ induced by Ψ), and the pages of the open book are given by the (relative) fibre sum

of the original page along the nested pages (with respect to the isomorphism of $\nu\pi_i'^{-1}(\theta) \subset T\pi^{-1}(\theta)$ *induced by* Ψ*), i.e.* $\overline{\Pi^{-1}(\theta)} = \#_{\Psi|_{\nu(\Sigma_0')_\theta}} \Sigma_\theta$. □

In the following we are going to extract the remaining information to express $\#_\Psi M$ in terms of an abstract open book, that is we describe a recipe to find the monodromy. Let us take a closer look at the nested open books within the ambient manifold. Let X be a vector field transverse to the interior of the ambient pages, vanishing on the binding, and normalised by $\pi^* d\theta(X) = 1$. Recall from Section 2.1 that the time-2π map ϕ of the flow of X yields the monodromy of the ambient open book. Furthermore, if we assume that X is tangent to the submanifolds M_i', we obtain abstract nested open book descriptions (Σ_i, ϕ_i) of M_i' within the abstract ambient open book (Σ, ϕ). Moreover, by adapting the vector field if necessary, we can choose embeddings of the normal bundles of M_i' such that the fibres are preserved under the flow of X. The normal bundles of M_0' and M_1' being isomorphic translates into the condition that the normal bundles $\nu\Sigma_i'$ of the induced (abstract) nested pages in the ambient (abstract) page Σ are ϕ-equivariantly isomorphic. Note that we have not specified such an isomorphism yet. A natural choice seems to be the isomorphism induced by Ψ, to be more precise, the restriction Ψ_0 of the isomorphism Ψ to $N_0 \cap \Sigma_{\theta=0}$. However we will see below that this is not the right choice in general, i.e. the abstract description of the open book in Lemma 3.3 does not equal $(\#_{\Psi_0}\Sigma, \phi)$ in general. It turns out we have to adapt the monodromy and add the corresponding *twist map*, which will be described in the following.

For the remaining part of the section we identify $\nu M_i'$ with the quotient

$$\big(\nu\Sigma_i' \times [0, 2\pi]\big)/\sim_\phi .$$

Now let Ψ_0 be the ϕ-equivariant fibre-orientation reversing isomorphism of $\nu\Sigma_i'$ within $T\Sigma$ induced by the restriction of Ψ. Moreover we define

$$\Psi_t := \Psi|_{\nu\Sigma_0' \times \{t\}}.$$

Note that each Ψ_t is isotopic to Ψ_0, the whole family $\{\Psi_t\}_t$ however defines an (*a priori*) non-trivial loop of maps $\nu\Sigma_0' \to \nu\Sigma_1'$ based at Ψ_0. By choosing suitable bundle metrics, this loop yields an (*a priori*) non-trivial loop $\{\mathcal{D}_t\}_t$ of maps $\Sigma' \to S^1$ based at the identity via

$$\mathcal{D}_t(x) \cdot \Psi_0(q) := \Psi_t(q),$$

for $x \in \Sigma'$ and $q \neq 0$ a non-trivial point in the normal-fibre over x. With this in hand we can define a monodromy-like map of $\nu\Sigma_1'$ which is the identity in a neighbourhood of the zero section and outside the unit-disc bundle by

$$\mathcal{D}(q) := \mathcal{D}_{\boldsymbol{r}(x)} \cdot q,$$

where $\boldsymbol{r}$ is a radial cut-off function in the fibre which is 1 on the zero section and vanishes away from it. We call it the **twist map** induced by ϕ and Ψ. Given this map we can now give an abstract description of the open book in Lemma 3.3. Recall that we already identified the page as the fibre sum of the original page along the nested pages.

Lemma 3.4. *Let* Ψ_0*,* ϕ *and* $\mathcal{D}$ *be the maps described in the above paragraph (i.e.* ϕ *is the monodromy of* $M = (B, \pi)$ *adapted to the nested open books,* $\mathcal{D}$ *is the derived twist-map, and* Ψ_0 *is the restriction of* Ψ *to the normal bundle of the subpage at angle zero). Then the monodromy of the open book in Lemma 3.3 is given by* $\phi \circ \mathcal{D}$*, and the page is* $\#_{\Psi_0}\Sigma$. □

4. The push-off

In this section we describe a *push-off* of the binding of an open book which realises it as a nested open book. The push-off construction will enable us to describe a natural open book structure on the fibre connected sum of two open books along their diffeomorphic bindings. The notion *push-off* may be a bit misleading, as the result of our construction is not a push-off in the ordinary sense but only close to it. In particular, the binding and its push-off will intersect. We will first describe how the binding is being pushed away from itself and then introduce a natural framing of the pushed-off copy in Subsection 4.1, which will be equivalent to the canonical *page framing* of the binding. In Subsection 4.2 we show that the push-off can be realised as an *abstract nested open book*.



Figure 2. The page Σ'_θ pushed into Σ_θ

Let M be a manifold with open book decomposition (Σ, ϕ) and binding B which also admits an open book decomposition (Σ', ϕ'). We denote the fibration maps by $\pi\colon M \setminus B \to S^1$ and $\pi'\colon B \setminus B' \to S^1$, respectively. Our aim is to define a push-off B^+ of the binding B in such a way that each page Σ'_θ of the binding open book is pushed into Σ_θ, the page corresponding to the same angle θ in the ambient open book. As all our constructions are local in a neighbourhood of the binding B, we can assume, without loss of generality, that ϕ is the identity.

Identify a neighbourhood of the binding $B' \subset B$ of the open book of the binding B with $B' \times D^2$ with coordinates (b', r', θ') such that (r', θ') are polar coordinates on the D^2-factor and θ' corresponds to the fibration π' – these are standard coordinates for a neighbourhood of a binding of an open book. We will also use Cartesian coordinates x', y' on the D^2-factor. Analogously, we have coordinates (b, r, θ) in a neighbourhood of $B \subset M$ with the corresponding properties. Combining these, we get two sets of coordinates on $(B' \times D^2) \times D^2 \subset M$:

$$(b', r', \theta', r, \theta) \quad \text{and} \quad (b', x', y', x, y).$$

First, we will describe the geometric idea of the push-off by considering just a single page Σ_θ of the open book before defining it rigorously afterwards, see Figure 2. The page Σ' of the binding open book is pushed into the page Σ. The push-off depends

on the radial direction r' only and is invariant in the B'-component. In particular, the boundary of the page Σ' stays fixed. We divide the collar neighbourhood in Σ' into four parts by the collar parameter r'. The outermost one consisting of points in Σ' with $r' \leq \epsilon_1$ is mapped to run straight into the r-direction of the ambient page Σ. The innermost part consisting of points with $r' \geq \epsilon_3$ is translated by a constant c into the r-direction. This translation is extended over the whole of Σ'. On the rest of the collar the push-off is an interpolation between these innermost and outermost parts. This is done such that points with $\epsilon_1 \leq r' \leq \epsilon_2$ are used to interpolate in r-direction and points with $\epsilon_2 \leq r' \leq \epsilon_3$ in r'-direction.

Let $f, h\colon \mathbb{R}_0^+ \to \mathbb{R}$ be the smooth functions described in Figure 3. Recall that B can be decomposed as $(B' \times D^2) \cup \Sigma'(\phi')$. Let $g\colon B \to B \times D^2 \subset M$ be the embedding defined by

$$g(b) = \begin{cases} \big((b', f(r') \cdot e^{i\theta'}), h(r') \cdot e^{i\theta'}\big) & \text{for } b = (b', r'e^{i\theta'}) \in B' \times D^2 \\ \big([x', \theta'], c \cdot e^{i}\theta'\big) & \text{for } b = [x', \theta'] \in \Sigma'(\phi'). \end{cases}$$

Observe that g is well-defined and a smooth embedding.

Definition 4.1. *We define the* ***push-off*** B^+ ***of*** B *as the image of the embedding g defined above, i.e. we define*

$$B^+ := g(B).$$

Observe that we can easily obtain an isotopy between the binding B and the push-off B^+ by parametrising f and h.



FIGURE 3. The functions f and h

Remark 4.2. We call the submanifold B^+ a *push-off* of B although it is not a push-off in the usual sense since B and B^+ are not disjoint. However, it generalizes the notion of a push-off of a *transverse* knot in a contact manifold. Note that $B^+ \cap \Sigma_\theta \cap \{r = c\}$ is a copy of the interior of the page Σ'_θ of the binding and $B^+ \cap \Sigma_\theta \cap \{r = r_0 < c\} \cong B' \times \{r_0\}$.

4.1. **Framings.** The fibre connected sum explained in Section 2.2 requires the submanifolds to have isomorphic normal bundles and explicitly uses a given bundle isomorphism. The binding of an open book has trivial normal bundle. Hence it is sufficient to specify a framing to be able to perform a fibre connected sum along the binding. In this section we will discuss a natural framing of the binding of an open book and introduce a corresponding framing for its push-off.

Let $N \subset M$ be a submanifold with trivial normal bundle. A *framing* of N is a trivialisation of its normal bundle. If the codimension of N in M equals two, we can consider the normal bundle as a complex line bundle, which can be trivialised by a nowhere-vanishing section. Thus, a framing of a codimension two submanifold with trivial normal bundle can be given by specifying a push-off, or equivalently a non-zero vector field along the submanifold that is nowhere tangent. We call two framed submanifolds equivalent if they are isotopic through framed submanifolds. Note that a framing of the binding B can also be specified by a homotopy class of maps from B to S^1. If B is simply-connected, such a map lifts to the universal cover $\mathbb{R}$ and thus is null-homotopic, i.e. the framing is unique.

A natural framing of the binding $B \subset M$ of an open book is the **page framing** obtained by pushing B into one fixed page of the open book. We denote the page framing given by $-\partial_x$ by F_0, i.e.

$$F_0 := -\partial_x.$$

Next we are going to define a framing for the push-off B^+. Let $\widetilde{u}\colon M \to \mathbb{R}$ be a smooth function such that

- $\widetilde{u} \equiv 0$ near B and on $B' \times \{r' \leq \epsilon\} \times D^2_{c-\epsilon}$,
- $\widetilde{u} \equiv 1$ on $B' \times \{r' \geq \epsilon\} \times \{r = c\}$ and outside $\{r \leq c + \epsilon\}$,
- $\widetilde{u}$ is monotone in r'- and r-direction.

With this in hand we define a framing of the push-off B^+ by

$$F_1 := -(1 - \widetilde{u})\partial_{x'} - \widetilde{u} \cdot (\sin^2\theta \partial_{x'} - \cos\theta \partial_r).$$

One easily checks that this is indeed nowhere tangent to B^+. We now show that the push-off B^+ with the framing F_1 is equivalent to the binding B with its natural page framing F_0.

Lemma 4.3. *The framed submanifolds (B, F_0) and (B^+, F_1) are isotopic.*

Proof. The strategy of the proof is as follows. We use an *intermediate push-off* $\widetilde{B}$ which is smooth outside a singular set and consider the framings F_0 and F_1 as framings on this intermediate push-off. Here, *framing* means an honest framing on the smooth part that extends continuously. We then describe a homotopy of the framings over $\widetilde{B}$ which has the property that it still defines a framing when the intermediate push-off is smoothened (cf. Definition und Notiz (13.12) in [2]). This then yields the desired isotopy of the framed submanifolds (B, F_0) and (B^+, F_1).

Let c be the constant as in the definition of the push-off B^+. Then we define the *intermediate push-off* $\widetilde{B}$ as the image of $\widetilde{g}\colon B \to B \times D^2 \subset M$ defined by

$$\widetilde{g}(b) := \begin{cases} \big(b, r'(b), \pi'(b)\big) & \text{for } r' \leq c, \\ \big(b, c, \pi'(b)\big) & \text{for } r' \geq c. \end{cases}$$

Note that the intermediate push-off has singular points in $r = c$. On the intermediate push-off the framings (which we continue to denote by F_0 and F_1 and leave out the base point for ease of notation) are as follows: The page framing is given by $F_0 = \partial_x$ or, written in polar coordinates,

$$F_0 = \cos\theta \partial_r - \sin\theta \partial_\theta,$$

with $\theta = \pi(\widetilde{g}(b))$. The second framing, the framing of the push-off, is given by

$$F_1 = t\cos\theta' \partial_r - (1 - t + t\sin^2\theta')\cos\theta' \partial_{r'} + (1 - t + t\sin^2\theta')\sin\theta' \partial_{\theta'}.$$

Here $t = t(b)$ is a function interpolating between 0 and 1 such that

- t only depends on the r'-direction,
- $t = 0$ for $r' \leq c - \epsilon$,
- $t = \frac{1}{2}$ for $r' = c$,
- $t = 1$ for $r' \geq c + \epsilon$.

We define

$$F_h := (1-h)F_0 + hF_1$$

and show that F_h defines a framing for every $h \in [0,1]$. The remainder of the proof is a mere calculation. The tangent space of $\widetilde{B}$ is defined outside of the set $\{r = c\}$ and is spanned by TB' and, depending on the value of t, the vectors

- $\partial_r + \partial_{r'}$ and $\partial_\theta + \partial_{\theta'}$, where $t < \frac{1}{2}$, or
- $\partial_{r'}$ and $\partial_\theta + \partial_{\theta'}$, where $t > \frac{1}{2}$.

Omitting the B'-direction and using vector notation in the ordered basis $\partial_r, \partial_{r'}, \partial_\theta$, $\partial_{\theta'}$, we have

$$F_h = \begin{pmatrix} \cos\theta(1-h+ht) \\ -\cos\theta h(1+t\sin^2\theta' - t) \\ -(1-h)\sin\theta \\ h(1+t\sin^2\theta' - t)\sin\theta' \end{pmatrix}.$$

We consider the cases $t \geq \frac{1}{2}$ and $t \leq \frac{1}{2}$ separately.

Case 1: $t \geq \frac{1}{2}$. In the chosen basis the tangent space to $\widetilde{B}$ consists of vectors of the form

$$\big(0,\, a,\, b,\, b\big)^\intercal$$

with $a, b \in \mathbb{R}$. Assume that $\cos\theta = 0$. Then the framing F_h becomes

$$\big(0,\, 0,\, -(1-h)\sin\theta,\, h(1+t\sin^2\theta' - t)\sin\theta'\big)^\intercal$$

and we have $\sin\theta \neq 0$. This vector being tangent to $\widetilde{B}$ requires $-1 = ht(\sin^2\theta' - 1)$, a contradiction since $0 \leq ht \leq 1$ and $(\sin^2\theta' - 1) \in (-1, 0]$. Otherwise, i.e. if $\cos\theta \neq 0$, the tangency condition yields $(1 - h + ht) = 0$, but this equation does not have a solution for $t \geq \frac{1}{2}$.

Case 2: $t \leq \frac{1}{2}$. Here the tangent space to $\widetilde{B}$ consists of vectors of the form

$$\big(a,\, a,\, b,\, b\big)^\intercal$$

with $a, b \in \mathbb{R}$. By the argument above we can rule out the case $\sin\theta \neq 0$. For $\sin\theta = 0$, we have $\cos\theta \neq 0$ and F_h equal to

$$\big(\cos\theta(1-h+ht),\, -\cos\theta h(1+t\sin^2\theta' - t),\, 0,\, 0\big)^\intercal.$$

Thus the tangency condition leads to $1 = -h + ht + h - ht = 0$, a contradiction.

It follows that F_h is never of the form

$$\big((1-t)a,\, a,\, b,\, b\big)^\intercal$$

and hence does indeed define a framing on a smoothened intermediate push-off. It follows that (B, F_0), $(\widetilde{B}, F_0)$, $(\widetilde{B}, F_1)$ and (B^+, F_1) are isotopic. □

4.2. **The push-off as an abstract nested open book.** The push-off B^+ is clearly an embedded nested open book of $M = (B, \pi)$. The aim of this section is to obtain a description of the push-off as an *abstract* nested open book, and ultimately as a *framed* abstract nested open book by altering the monodromy of the abstract open book (Σ, ϕ). To this end, the monodromy of the binding is used to define a vector field X' on $B \setminus B'$, which can then be extended to a neighbourhood of B in M. By cutting off this vector field appropriately, we alter the vector field X on $M \setminus B$ corresponding to the monodromy ϕ to get a new monodromy of the form $\phi \circ \psi$. This will realise the push-off as an abstract nested open book. Finally, the monodromy of the ambient open book is changed again in such a way that the framing of a page of the push-off is independent of the ambient page. This means that the framed push-off can then be regarded as a framed abstract nested open book.

Identify a neighbourhood of $B' \subset B$ with $B' \times D^2$ as above, i.e. the pages are defined by the angular coordinate θ'. Also denoting the coordinate on S^1 by θ', we can define a non-vanishing 1-form on $B \setminus B'$ by the pull–back of $d\theta'$ under the fibration map $\pi' \colon B \setminus B' \to S^1$. With the help of this 1-form we can extend the vector field $r^2\partial_{\theta'}$ to a vector field X' on B by prescribing the condition $(\pi')^* d\theta'(X') = 1$. The vector field X' can furthermore be extended trivially to a neighbourhood $B \times D^2$ of B in M. We obtain an abstract open book description of M by regarding the time-2π map of a certain vector field on $M \setminus B$ which satisfies conditions analogue to the ones used in the construction of X' above. Let X denote the vector field that recovers the abstract open book (Σ, ϕ). Let $u \colon \mathbb{R}_0^+ \to \mathbb{R}$ be the smooth function depicted in Figure 4 with c as in the definition of the push-off B^+. Then $\widetilde{X} := X + uX'$ on $M \setminus B$ defines a vector field on $M \setminus B$.



FIGURE 4. The function u

We claim that $\widetilde{X}$ realises the push-off B^+ as an abstract nested open book of an abstract open book description of M. Observe that, as X' is tangent to the pages of (B, π), the condition $\pi^* d\theta(\widetilde{X}) = 1$ is satisfied and that $\widetilde{X}$ and X coincide near the binding B. Thus, the vector field $\widetilde{X}$ does indeed yield an abstract open book description of M. Furthermore, the vector field $\widetilde{X}$ is tangent to the push-off B^+, which means that it realises B^+ as an abstract nested open book of the abstract ambient open book. The monodromy is given by the time-2π flow of $\widetilde{X}$. However, we want to give a description that better encodes the change of the monodromy ϕ of the ambient open book we started with in terms of the monodromy of the binding.

We denote the flow of X' on B by ϕ'_t and use it to define diffeomorphisms ψ_t of a neighbourhood $B \times D^2$ of B in M:

$$\psi_t(b, r, \theta) = (\phi'_t(b), r, \theta).$$

Definition 4.4. *Define a diffeomorphism $\psi\colon M \to M$ via $\psi := \psi_{2\pi u(r)}$ and refer to it as **Chinese burn**[2] along B. By abuse of notation, its restriction to a single page Σ_θ is also denoted by ψ.*



FIGURE 5. A Chinese burn along a boundary component.

Observe that the monodromy of the abstract open book obtained from the vector field $\widetilde{X}$ is $\phi \circ \psi$, i.e. the push-off B^+ yields an an abstract nested open book of $(\Sigma, \phi \circ \psi)$. We thus proved the following statement.

Lemma 4.5. *The push-off B^+ induces an abstract nested open book of $(\Sigma, \phi \circ \psi)$ with page diffeomorphic to Σ' (more concretely, the page is $g|_{\Sigma'_0}(\Sigma'_0) \cong \Sigma'$), where ψ is a Chinese burn along B.*

We constructed the push-off B^+ inside the manifold $M(\Sigma, \phi)$ and equipped it with a natural framing F_1 corresponding to the page framing. In particular, the push-off is a *framed* nested open book, i.e. a nested open book with a specified framing. The previous lemma shows that the push-off also defines an *abstract* nested open book of $(\Sigma, \phi \circ \psi)$. However, the framing F_1 does *a priori* not give a framing in the abstract setting since it is not invariant under the monodromy. We call an abstract nested open book with a framing which is invariant under the monodromy a **framed abstract nested open book**.

Remark 4.6. Given two diffeomorphic framed nested open books with pages $\Sigma'_0, \Sigma'_1 \subset (\Sigma, \phi)$ and isotopic normal bundles, it is easy to obtain an open book structure of their fibre connected sum. The new page is

$$\widetilde{\Sigma} := \big(\Sigma \setminus (\nu\Sigma'_0 \cup \nu\Sigma'_1)\big)/\sim$$

with the identification induced by the given framings, and the old monodromy ϕ restricts to the new monodromy.

[2]In common speech, a *Chinese burn* is the "act of placing both hands on a person's arm and then twisting it to produce a burning sensation" (cf. *oxforddictionaries.com*).

A natural framing of the push-off in the abstract setting is the following: We define the **constant framing** F_2 as

$$F_2 \colon \widetilde{u}\partial_r - (1-\widetilde{u})\partial_{x'},$$

where $\widetilde{u}$ is the restriction of the function $\widetilde{u} \colon M \to \mathbb{R}$ defined in Section 4.1 to a page Σ, i.e. it is smooth with the following properties:

- $\widetilde{u} \equiv 0$ near B and on $B' \times \{r' \leq \epsilon\} \times D^2_{c-\epsilon}$,
- $\widetilde{u} \equiv 1$ on $B' \times \{r' \geq \epsilon\} \times \{r = c\}$ and outside $\{r \leq c + \epsilon\}$,
- $\widetilde{u}$ is monotone in r'- and r-direction.

To realise the push-off as a framed abstract nested open book with framing F_2, we have to alter the monodromy of the ambient abstract open book. We will change the monodromy by a certain diffeomorphism of the page fixing the push-off, the so-called *twist map*. Let $\sigma \colon \mathbb{R} \to \mathbb{R}$ be a cut-off function with $\sigma(0) = 0$ and $\sigma(\epsilon) = 1$ and

$$\tau_{\pm}^{\sigma(r)} \colon D^2 \to D^2$$

a smoothened Dehn twist of the disc. That is, a diffeomorphism with the qualitative behaviour of

$$(s, \theta) \mapsto (s, \theta \pm 2\pi\sigma(r)(1-s)),$$

smoothened near the boundary and the origin, such that the origin is an isolated fixed point and a neighbourhood of the boundary is fixed. This can be achieved by constructing it as the flow of an appropriate vector field. Recall that, by construction, the intersection of the push-off of the binding B with a single page Σ of the ambient open book is a copy of a page Σ' of the open book of the binding. In particular, we can identify a tubular neighbourhood of $B^+ \cap \Sigma_0$ in Σ_0 with $\Sigma' \times D^2 \subset \Sigma$. Observe that the r-coordinate can be regarded as a collar parameter on Σ'. We define a diffeomorphism of a neighbourhood of the collar by

$$\begin{aligned} \partial\Sigma' \times [0,\epsilon] \times D^2 &\to \partial\Sigma' \times [0,\epsilon] \times D^2 \\ (b', r, p) &\mapsto (b', r, \tau_-^{\sigma(r)}(p)). \end{aligned}$$

We can extend this to a diffeomorphism $\Sigma' \times D^2 \to \Sigma' \times D^2$ of the whole tubular neighbourhood of Σ' via $\mathrm{id}_{\Sigma'} \times \tau_-^1$. Furthermore, this map can be extended to a self-diffeomorphism $\mathcal{D} \colon \Sigma \to \Sigma$ of the page Σ via the identity.

Definition 4.7. *The diffeomorphism $\mathcal{D} \colon \Sigma \to \Sigma$ is called **twist map**.*

The twist map $\mathcal{D}$ is isotopic to the identity, so we have $M_{(\Sigma,\phi\circ\psi)} \cong M_{(\Sigma,\phi\circ\psi\circ\mathcal{D})}$.

Lemma 4.8. *The push-off B^+ with its induced framing F_1 corresponds to the framed abstract nested open book of $(\Sigma, \phi \circ \psi \circ \mathcal{D})$ with page $g|_{\Sigma'_0}(\Sigma'_0) \cong \Sigma'$ framed by the natural framing F_2.*

Proof. The push-off B^+ was constructed in $M_{(\Sigma,\phi)}$. To prove the lemma, we will mostly work in $M_{(\Sigma,\phi\circ\psi)}$. We begin by describing the push-off B^+ seen in this manifold. A diffeomorphism between $M_{(\Sigma,\phi)}$ and $M_{(\Sigma,\phi\circ\psi)}$ is given by the map $\Theta \colon M_{(\Sigma,\phi)} \to M_{(\Sigma,\phi\circ\psi)}$ defined by

$$\Theta(p) = \psi_{-\pi(p)u(\mathbf{r}(p))}(p),$$

where $\mathbf{r} \colon \Sigma \to \mathbb{R}$ is a function that coincides with the collar parameter r near B and is constantly extended to all of Σ. In particular, it preserves the pages Σ_θ

and the r-direction, and is induced by the inverse of the time-θ map of the binding monodromy on the $\{r = c\}$-slice in Σ_θ. We denote the image of B^+ under the diffeomorphism Θ by B_1. The diffeomorphism Θ also transports the framing F_1 of B^+ to a framing of B_1, that we also denote by F_1.

On the other hand, the embedded nested open book B_1 induces the abstract nested open book from Lemma 4.5. Hence, as the twist map $\mathcal{D}$ fixes its pages, the framing F_2 also induces a framing of B_1 by considering the diffeomorphism from $M_{(\Sigma,\phi\circ\psi\circ\mathcal{D})}$ to $M_{(\Sigma,\phi\circ\psi)}$ coming from the twist map $\mathcal{D}$ (cf. Figure 6).

It remains to show the equivalence of the two framings of B_1. Note that $\Theta\colon M_{(\Sigma,\phi)} \to M_{(\Sigma,\phi\circ\psi)}$ leaves the ∂_r-direction of the framing unchanged and is a rotation in the B-directions. More concretely, we have

$$\partial_{x'} \mapsto \cos(-\theta' u(h_s(r')))\partial_{x'} + \sin(-\theta' u(h_s(r')))\partial_{y'}.$$

Thus, the framing F_1 of B_1 obtained through the diffeomorphism is

$$b \mapsto \left(\widetilde{b},\ \widetilde{u}\left(-\sin^2\theta'\left(\lambda_1\partial_{x'} + \lambda_2\partial_{y'}\right) + \cos\theta'\partial_r\right) + (1-\widetilde{u})\left(-\lambda_1\partial_{x'} + \lambda_2\partial_{y'}\right)\right)$$

with

$$\widetilde{b} = \psi_{-\theta\cdot u(\mathbf{r}(g_s(b)))}\left(g_s(b)\right)$$

and

$$\lambda_1 = \cos(-\theta' u(h_s(r'))),\ \ \lambda_2 = \sin(-\theta' u(h_s(r'))).$$

The framing F_2 of B_1 in $M_{(\Sigma,\phi\circ\psi\circ\mathcal{D})}$ is mapped to

$$\widetilde{u}(\cos\theta'\partial_r + \sin\theta'\partial_{y'}) + (1-\widetilde{u})\cdot(-1)\cdot(\cos(-\theta' d(r))\partial_{x'} + \sin(-\theta' d(r))\partial_{y'})$$

under the diffeomorphism induced by the twist map $\mathcal{D}$ from $M_{(\Sigma,\phi\circ\psi\circ\mathcal{D})}$ to $M_{(\Sigma,\phi\circ\psi)}$. We can choose $d(r) = u(c_s)$ in the definition of $\mathcal{D}$ and simply interpolate between $\mathcal{D}(F_2)$ and F_1, where a little care is needed only for $\theta = \frac{\pi}{2}$ and $\theta = \frac{3\pi}{2}$. □



FIGURE 6. The framing F_2 and the twist $\mathcal{D}$

5. Proof of Theorem 1.1

We are now able to give an explicit open book decomposition for the binding sum operation and thus prove our main result. The proof is divided into two parts. First, we show the existence of a natural open book decomposition of the binding sum and specify its monodromy in the abstract setting. In the second part we prove that the resulting page can be obtained from the original pages by two consecutive generalised 1-handle attachments.

5.1. **Proof of the 1st part: Existence and identifying the monodromy.** Let M be a manifold with open book fibration (B, π) two of whose binding components, denoted by B_0 and B_1, admit diffeomorphic open books. By Lemma 4.3 B_0 and B_1 are isotopic to their respective push-offs B_i^+. Hence, the binding sum along the B_i (with respect to the page framing) can be replaced by the fibre connect sum along the push-offs B_i^+ with the framing F_1. Since the push-offs are nested open books $(B_i^+, \pi_i := \pi|_{B_i^+})$ of (B, π), the fibre connected sum carries a natural open book structure coinciding with the original open book fibration away from the push-offs by Lemma 3.3. A fibre of this new open book fibration is the fibre connected sum of the corresponding original fibres along the fibres of the nested open books.

In the abstract setting we have $M = M_{(\Sigma,\phi)}$ and the binding components B_0 and B_1 admit an abstract open book decomposition (Σ', ϕ'). By Lemma 4.8, the push-offs B_i^+ induce framed nested open books of $(\Sigma, \phi \circ \psi \circ \mathcal{D})$, and we have $M = M_{(\Sigma,\phi)} \cong M_{(\Sigma,\phi\circ\psi\circ\mathcal{D})}$. Here, ψ denotes the Chinese burn along the relevant binding components B_i (see Definition 4.4) and $\mathcal{D}$ the corresponding twist map (see Definition 4.7). As described in Remark 4.6, the abstract open book description of the fibre connected sum follows easily because the framing of a framed nested open book is compatible with the ambient monodromy. The new page is

$$\widetilde{\Sigma} := \big(\Sigma \setminus (\nu\Sigma_0' \cup \nu\Sigma_1')\big)/\sim$$

with the identification induced by the given framings, i.e. it is the fibre connected sum of the original page Σ along the framed nested pages Σ_i' of the framed nested open books of $(\Sigma, \phi\circ\psi\circ\mathcal{D})$ corresponding to the B_i, and the monodromy is induced by $\phi \circ \psi \circ \mathcal{D}$. □

5.2. **Proof of the 2nd part: Handle interpretation of the page.** In this section we explain how the binding connected sum of two open books with diffeomorphic bindings can be regarded as two consecutive generalised 1-handle attachments. Let Σ be the ambient page, i.e. $\Sigma \cong \overline{\pi^{-1}(0)}$ with (B, π) the open book, and let $B_0, B_1 \subset B = \partial\Sigma$ be two diffeomorphic binding components with common open book. We already discussed what the open book for the binding sum looks like. For ease of notation, let Σ_i' denote the subpage of the push-off B_i^+ of B_i as well as the page of B_i corresponding to a fixed angle, zero say, for $i = 0, 1$. Let $\nu\Sigma_i' \subset \Sigma$ be their embedded normal bundles. Then the new page is given by

$$\widetilde{\Sigma} := \overline{\Pi^{-1}(0)} = \#_{\Psi|_{\nu\Sigma_0'}}\Sigma.$$

Equivalently, in the abstract setting, we can identify $\nu\Sigma_i'$ with $\Sigma_i' \times \mathrm{int}D^2 \subset \Sigma_i' \times \mathbb{C}$ via the constant framing of the constant push-offs and obtain

$$\widetilde{\Sigma} := \overline{\Pi^{-1}(0)} = \Big(\Sigma \setminus \big(\nu\Sigma_0' \cup \nu\Sigma_1'\big)\Big)/(x, z) \sim (x, \bar{z}).$$

Note that $\Sigma_i' \times \{1\} \subset \nu\Sigma_0'$ is isotopic in $\Sigma \setminus \nu\Sigma_0'$ to the corresponding page $\Sigma_i' \subset B_i \subset \partial\Sigma$ of the respective binding component B_i. Let $\Sigma' \times [-1, 0]$ and $\Sigma' \times [0, 1]$ denote the traces of the obvious isotopies fixing $\partial\Sigma' \times \{1\}$. Observe furthermore that these two isotopies glue together in $\overline{\Pi^{-1}(0)}$ to an isotopy whose image we denote by $\Sigma' \times [-1, 1]$ (cf. Figure 7), which clearly is the co-core of a generalised 1-handle $H^{(2)}_{\Sigma'}$.

If we cut open the new page $\widetilde{\Sigma}$ along the co-core of this handle and consider the effect in the original page Σ, the areas $\Sigma_i' \times \partial D^2$ of identification become



FIGURE 7. The dark grey region on each side is the neighbourhood of a single page of the push-off of binding components. The light grey region defines the co-core of a generalised 1-handle $H^{(2)}_{\Sigma'}$. After cutting along the co-core of $H^{(2)}_{\Sigma'}$, the dark grey region descends to the co-core of another generalized 1-handle $H^{(1)}_{\Sigma'}$.

diffeomorphic to $\Sigma_i' \times [0, 2\pi]$ since $\Sigma_i' \times \{1\}$ is removed. Hence, we end up with a single embedded copy of $\Sigma' \times [0, 2\pi]$ in the quotient $\widetilde{\Sigma}$. This is the co-core of another generalised 1-handle $H^{(1)}_{\Sigma'}$. Cutting along $H^{(1)}_{\Sigma'}$ then gives back the original page Σ (cf. Figure 7). Thus, we have

$$\overline{\Pi^{-1}(0)} = \left(\Sigma \sqcup H^{(1)}_{\Sigma'} \sqcup H^{(2)}_{\Sigma'}\right) / \sim,$$

where we attach the first handle $H^{(1)}_{\Sigma'}$ along two copies of Σ', i.e. along one page of each of the two binding components B_0 and B_1. The second handle $H^{(2)}_{\Sigma'}$ is then attached to $D^1 \times \Sigma \times S^0 \subset \partial H^{(1)}_{\Sigma'}$ (see Figure 8). □

Example 5.1. Let $M = M_0 \sqcup M_1$ with M_i the four-dimensional sphere S^4 with open book decomposition $(\Sigma_i = D^3, \mathrm{id})$. Then the binding has two components B_i, both a 2-sphere with the unique open book decomposition $(\Sigma_i' = D^1, \mathrm{id})$. We have $M_i = D^3 \times S^1 \cup S^2 \times D^2$, so performing the binding sum on M along the B_i yields $S^3 \times S^1$. By Theorem 1.1, this has a natural open book decomposition obtained by forming the sum along the push-off of the binding.

Pushing a page $\Sigma_i' = D^1$ of the binding open book into the page $\Sigma_i = D^3$ and then removing a neighbourhood of Σ_i', turns Σ_i into a solid tours $S^1 \times D^2$. Hence, the page in the resulting open book decomposition of the binding sum consists of two solid tori glued together along a neighbourhood of $S^1 \times \{*\} \subset S^1 \times \partial D^2$, i.e. it is a solid torus. Note that we do not have to specify a framing since S^2 is simply-connected and thus possesses a unique framing.

Recall that the Chinese burn ψ is non-trivial only in a collar neighbourhood of the binding. To identify the new monodromy it is helpful to imagine the collar as split into two parts. The Chinese burn twists the outer part of the collar and then twists back on the inner part. Now observe that the new monodromy is isotopic to the identity. Indeed, the twist map $\mathcal{D}$ on the new page cancels with the Chinese burn on the outer part and the Chinese burn restricted to the inner part is isotopic



FIGURE 8. The dark and light grey region define cores of generalized 1-handles $H^{(1)}_{\Sigma'}$ and $H^{(2)}_{\Sigma'}$ respectively. Note that, in the particular case we pictured here, the 2-fold handle attachment yields a solid torus $S^1 \times D^2$.

to the identity also on the new page (see also the 3-dimensional case sketched at the end of this example).

Alternatively, we can also construct the new page by two consecutive generalised 1-handle attachments. In the present situation a generalised 1-handle $H_{\Sigma'}$ is just $D^1 \times D^1 \times D^1$. The first handle attachment described above turns $\Sigma = D^3 \sqcup D^3$ into a single ball, the second handle attachment results in a solid torus, see Figure 8.

It is also worth comparing this with the situation in one dimension lower as discussed in [10], i.e. the binding sum of two copies of (D^2, id). with page framing (here, a framing has to be specified). Then the new page is an annulus and the monodromy is isotopic to the identity: the Chinese burn part of the monodromy are two trivial negative Dehn twists and two non-trivial positive Dehn twists which cancel with the two positive Dehn twists forming the twist map (see [10, Theorem 3]).

6. Application to contact topology

In this section we want to show that the binding sum of contact open books yields an open book and a contact structure that again fit nicely together. For an introduction to open books in contact topology we kindly refer the reader to [5, 6, 16].

A positive contact structure ξ on an oriented manifold M is **supported** by an open book structure (B, π) if it can be written as the kernel of a contact form α inducing a positive contact structure on B and such that $d\alpha$ induces a positive symplectic structure on the fibres of π. Such a contact form α is then called **adapted** to the open book. Note that a contact form is adapted to an open book (B, π), where the binding B is a contact submanifold, if and only if its Reeb vector field is positively transverse to the fibres of π (cf. [16, Lemma 2.13]). The flow of the Reeb vector field R_α of an adapted contact form does not necessarily preserve the open book structure (i.e. maps pages to pages). However, after multiplying R_α with an accordingly defined positive function h on M, the flow of the new vector

field hR_α does. We can realize hR_α as the Reeb vector field $R_{\alpha'}$ of another contact form α' for ξ. We call $R_{\alpha'}$ an **adapted** Reeb vector field.

Now let $(M, \xi = \ker \alpha)$ be a contact manifold supported by an open book (B, π). Let X be a vector field on M such that

- it is transverse to the pages and satisfies $d\theta(X) = 1$ on $M \setminus B$,
- the restriction of $\mathcal{L}_X d\alpha$ to any page vanishes,
- it equals ∂_θ on a neighbourhood $B \times D^2 \subset M$ of the binding, where (r, θ) are polar coordinates on the D^2-factor, and the open book fibration is given by the angular coordinate on D^2.

We call such a vector field X a **monodromy vector field**.

Given a monodromy field we get an associated abstract open book description of (M, ξ), and in turn an identification of (M, ξ) as a generalized Thurston-Winkelnkemper construction. In particular, such a vector field always exists if the open book and contact structure comes from a generalized Thurston-Winkelnkemper construction. Since, according to [3, Theorem 3.1.22], any contact manifold supported by an open book can be realised by a generalised Thurston-Winkelnkemper construction, we can always assume the existence of a compatible monodromy vector field. An adapted Reeb vector field is not a monodromy vector field as it does not fix the binding point-wise.

Recall that a **Lutz pair** h_1, h_2 is a pair of smooth functions $h_1 \colon [0,1] \to \mathbb{R}^+$ and $h_2 \colon [0,1] \to \mathbb{R}_0^+$ such that

- $h_1(0) = 1$ and h_2 vanishes like r^2 at $r = 0$,
- $h_1'(r) < 0$ and $h_2'(r) \geq 0$ for $r > 0$,
- all derivates of h_1 vanish at $r = 0$.

Note that the fibre connected sum construction can be adapted to the contact setting as follows.

Theorem 6.1 ([6, Theorem 7.4.3])**.** *Let (M, ξ) and (M', ξ') be contact manifolds of dimension $\dim M' = \dim M - 2$, where the contact structures ξ, ξ' are assumed to be cooriented; these cooriented contact structures induce orientations of M and M'. Let $j_0, j_1 \colon (M', \xi') \to (M, \xi)$ be disjoint contact embeddings that respect the coorientations, and such that there exists a fibre-orientation-reversing bundle isomorphism $\Phi \colon N_0 \to N_1$ of the normal bundles of $j_0(M')$ and $j_1(M')$. Then the fibre connected sum $\#_\Phi M$ admits a contact structure that coincides with ξ outside tubular neighbourhoods of the submanifolds $j_0(M')$ and $j_1(M')$.*

The binding of a contact open book decomposition is a contact submanifold and admits an open book structure. Furthermore, it has trivial normal bundle. Given two contact open books with contactomorphic bindings, we can thus perform the contact fibre connected sum along their bindings, and, topologically, also along the push-offs B_i^+ of the bindings. In fact, the latter also works in the contact world.

Lemma 6.2. *The push-off B^+ of the binding B of a contact open book (B, π) is a contact submanifold contact isotopic to the binding.*

Proof. We first show that the push-off B^+ of the binding B is a contact submanifold of (M, ξ). Since a collar neighbourhood of a fibre of π looks like the symplectisation of the binding B, we only have to prove this for a neighbourhood of the binding B' of the open book of B. On a neighbourhood $B' \times D^2 \times D^2$ (as described in

Section 4) we have a contact form

$$\alpha = \alpha'' + \frac{1}{2}r'^2 d\theta' + \frac{1}{2}r^2 d\theta,$$

where α'' is a contact form on B'. The push-off B^+, which arises as the image of the embedding $g\colon B \to M$, is, by definition, a contact submanifold of M if $g^*\alpha$ is a contact form on B. We compute

$$\begin{aligned} g^*\alpha \wedge (dg^*\alpha)^{n-1} &= (n-1)(ff' + hh')\alpha'' \wedge (d\alpha'')^{n-2} \wedge dr' \wedge d\theta' \\ &= (n-1)\frac{1}{r'}(ff' + hh')\alpha'' \wedge (d\alpha'')^{n-2} \wedge r' dr' \wedge d\theta'. \end{aligned}$$

For $r' \neq 0$ this is clearly positive because α'' is a contact form on B' and we have $(ff' + hh') > 0$ away from B'. Considering the limit

$$\lim_{r'\to 0} \frac{f(r')f'(r') + h(r')h'(r')}{r'} = \lim_{r'\to 0} \frac{h(r')h'(r')}{r'} = h'(0)^2 > 0$$

shows it is a volume form on all of $B' \times D^2$, which means that $g^*\alpha$ is a contact form. A similar calculation shows that the push-off B^+ is furthermore contact isotopic to the binding B. □

Lemma 6.3. *There is a contact form α on M and a neighbourhood U of B^+ in M with polar coordinates (s, ϑ) induced by the framing F_1, such that α is adapted to the contact open book structure (B, π) and its restriction to U is of the form*

$$\alpha_{|_U} = \alpha_{B^+} + s^2 d\vartheta,$$

where α_{B^+} is a contact form on the push-off B^+ of the binding B.

Proof. Assume that we have an adapted contact form α such that its Reeb vector field R is tangent to the push-off B^+. This can be obtained in exactly the same way as the monodromy vector field in the proof of Proposition 6.4. We can furthermore assume that we have

$$\alpha|_V = h_1 g_1 \alpha_{B'} + h_1 g_2 d\theta' + h_2 d\theta$$

for Lutz pairs h_1, h_2 depending on r and g_1, g_2 depending on r' in an open neighbourhood V of the nested binding B' in M.

The push-off B^+ is a contact submanifold by Lemma 6.2, i.e. $\alpha^+ := \alpha|_{TB^+}$ is a contact form on B^+. The framing F_1 of the push-off induces an identification of a neighbourhood of B^+ in M with $B^+ \times D^2$ in such a way that

$$(B^+ \times D^2) \cap \pi^{-1}(\theta) = (B^+ \cap \pi^{-1}(\theta)) \times D^2,$$

i.e. the normal directions to the push-off are contained in the pages. Then

$$\widetilde{\alpha} := \alpha^+ + s^2 d\vartheta$$

is a contact form on $B^+ \times D^2$. Here, (s, ϑ) denote polar coordinates on the D^2-factor. In the neighbourhood V of B' we can clearly choose s equal to $\sqrt{h_1} r'$ and $d\vartheta = d\theta'$. Our aim is to globalise this form by interpolating between its Reeb field $\widetilde{R}$ and the original Reeb field R in s-direction such that the interpolated vector field is positively transverse to the fibres of π and it is the Reeb vector field of a suitable contact form.

Observe that we have

$$\widetilde{R}|_{B^+} = R|_{B^+}, \tag{6.1}$$

so by continuity $\widetilde{R}$ is positively transverse to the pages in a neighbourhood of the push-off. On $B^+ \times D^2$, the original Reeb vector field R can be written as the contact vector field with respect to $\widetilde{\alpha}$ associated with the positive function $h := \widetilde{\alpha}(R)$. Now consider the likewise positive function $\widetilde{h} := (1-\eta)h + \eta$, where η is a smooth function on $B^+ \times D^2$ such that

- η is invariant in B^+- and ϑ-direction and monotone in s-direction,
- η equals 1 in a neighbourhood of $B^+ \times \{0\}$,
- η vanishes for $s \geq s_0$.

The contact vector field associated with $\widetilde{h}$ with respect to $\widetilde{\alpha}$ is

$$X = (1-\eta)R + \eta\widetilde{R} + (1-h)(X_\eta - \eta\widetilde{R}),$$

where X_η is the contact vector field (with respect to $\widetilde{\alpha}$) corresponding to the function η. Note that we have $d\eta(\widetilde{R}) = 0$. One then easily computes X_η to be

$$X_\eta = (\eta - \frac{s\eta'}{2})\widetilde{R} + \frac{\eta'}{2s}\partial_\vartheta$$

with $\eta' := \frac{\partial \eta}{\partial s}$.

Recall that we have $s = \sqrt{h_1}r'$ and $d\vartheta = d\theta'$ on the open neighbourhood V of B', and thus

$$\widetilde{\alpha}|_V = h_1 g_1 \alpha_{B'} + h_2 d\theta + s^2 d\vartheta = h_1 g_1 \alpha_{B'} + h_2 d\theta + h_1 g_2 d\theta' = \alpha|_V,$$

i.e. the two forms coincide on V. In particular, $h|_V = 1$ and $X|_V = R|_V$, so X is positively transverse to the fibres of π. It remains to inspect the situation outside of V. Both vector fields R and $\widetilde{R}$ are positively transverse to the pages. As $B^+ \setminus V$ is compact, the ∂_θ-component of $(1-\eta)R + \eta\widetilde{R}$ has a minimum m on $(B^+ \times D^2) \setminus V$. By taking s_0 small, we can achieve the product of $(1-h)\cdot(-\frac{s\eta'}{2})$ with the ∂_θ-component of $\widetilde{R}$ to be smaller than m since h is equal to 1 on B^+ by 6.1 and η can be chosen such that $|s\eta'|$ is bounded independently of the choice of s_0. Therefore, the vector field X is positively transverse on $(B^+ \times D^2) \setminus V$. Hence, X is positively transverse to the fibres of π on all of $B^+ \times D^2$.

As the function $\widetilde{h}$ is positive on $B^+ \times D^2$, we can rescale the contact form $\widetilde{\alpha}$ such that X is the Reeb vector field of the rescaled form. Notice that the form is adapted to the open book since X is positively transverse to the fibres of π. Outside the support of η, the vector field X coincides with the Reeb vector field R of the original (global) contact form α. We can thus glue the original form α to the rescaled form on $B^+ \times D^2$ to obtain the desired adapted contact form on M. Indeed, if we define U to be the interior of the set $\{\eta = 1\} \subset B^+ \times D^2$, this new contact form restricts to $\alpha_{B^+} + s^2 d\vartheta$ on U. □

6.1. **Proof of Theorem 1.3.** We can now prove the main theorem in the contact setting. The first part of the theorem is simply the statement of Lemma 6.2. For the proof of the adaptedness we will first treat the binding components B_0 and B_1 separately and will therefore drop the indices for ease of notation.

By Lemma 6.3, there is a contact form α on M that restricts to $\alpha_{B^+} + s^2 d\vartheta$ on a tubular neighbourhood $B^+ \times D^2$ of the push-off induced by the framing F_1. The contact form for ξ used in the contact fibre connected sum construction is of the form $\widetilde{\alpha} = \alpha_{B^+} + f(s)d\vartheta$ for an appropriate function f and coincides with α near $B^+ \times \partial D^2$ (see [6, proof of Theorem 7.4.3]). In particular, the Reeb vector

field of $\widetilde{\alpha}$ is still transverse to the fibres of π. Hence, the resulting contact form in the contact fibre connected sum construction of two contact open books along the push-offs of their contactomorphic bindings is again adapted to the resulting open book structure. □

6.2. **Recipe for an abstract description.** We conclude the paper with a recipe for an abstract description of the contact open book of the binding sum. We first want to construct a monodromy vector field which is tangent to the push-off B^+. The associated abstract open book of (M, ξ) then contains the push-off as an abstract nested open book.

Proposition 6.4. *The push-off can be realised as an abstract nested contact open book of an abstract contact open book for* (M, ξ).

Proof. We have a compatible contact form α on M which restricts to

$$\alpha = h_1\alpha' + h_2 d\theta$$

in a neighbourhood $B \times D^2$ of the binding. As usual, (r, θ) are polar coordinates on the D^2-factor, the open book fibration is given by the angular coordinate, (h_1, h_2) is a Lutz pair, and α' is a contact form on B. We can assume that α' is adapted to the open book structure on the binding B and furthermore that its Reeb field R' is an adapted Reeb vector field. We trivially extend this Reeb vector field to the neighbourhood $B \times D^2$ and still denote it by R'. Notice that this vector field is the contact vector field associated with the function h_1 and the vector field corresponding to h_2 is ∂_θ. Now consider the contact vector field

$$X = \partial_\theta + u \cdot R' + h_1(X_u - uR)$$

associated with the function $\widetilde{h} = h_2 + uh_1$, with u as in Section 4.2, X_u its contact vector field, and R the Reeb vector field of α. The latter computes as

$$R = \frac{1}{h_1 h_2' - h_1' h_2}(h_2' R' - h_1' \partial_\theta),$$

so we have $d\widetilde{h}(R) = 0$ and thus

$$\mathcal{L}_X d\alpha = 0.$$

Hence, X is clearly a monodromy vector field and it only remains to show that it is tangent to the push-off B^+.

We assumed the Reeb field R' of the binding to be an adapted Reeb field, i.e. it respects the pages of the open book structure of B. This means that $\partial_\theta + R'$ is tangent to the push-off B^+. Note that over $\{u = 1\}$ we have $X_u = R$ and hence $X = \partial_\theta + R'$. Therefore it remains to check the tangency condition where $0 < u < 1$. The tangency follows if X_u is tangent because both ∂_θ and R' are tangent to B^+ in that region (observe that for $r' = 0$, R' is tangent to B' thus tangent to B^+, and without any loss of generality we assume the complement of the innermost part of the push-off lies within $\{u = 1\}$). We can then compute X_u to be

$$X_u = uR - \frac{u' h_2}{h_1 h_2' - h_1' h_2} R' + \frac{u' h_1}{h_1 h_2' - h_1' h_2} \partial_\theta,$$

which is clearly tangent as well. □

In the following we are going to compute the symplectic monodromy of the open book described in Theorem 1.3 (or in Proposition 6.4, respectively). First, let us take a look at its page. Let $\nu_\varepsilon \Sigma_i'$ denote the complement of an ε-neighbourhood of the zero section of the normal bundle of the page Σ_i' of the nested open book within the ambient page Σ, where we identify Σ_i' and Σ with the closure of the fibres corresponding to angle zero in the respective open books. Furthermore, we can use the framing F_1 from the previous section to identify $\nu_\varepsilon \Sigma_i'$ with $\Sigma_i' \times (\varepsilon, 1) \times S^1$. Then the page is diffeomorphic to the quotient

$$\widetilde{\Sigma} := \overline{\Pi^{-1}(0)} = \Big(\Sigma \setminus \big(\nu\Sigma_0' \sqcup \nu\Sigma_1' \big) \sqcup \big(\Sigma' \times (-1,1) \times S^1 \big) \Big) / \sim,$$

where we identify with respect to the maps $j_i \colon \big(\Sigma_i' \times (\varepsilon, 1) \times S^1 \big) \to \big(\Sigma' \times (-1,1) \times S^1 \big)$ defined by

$$j_0(x', s, \vartheta) := (x', -s, -\vartheta) \quad \text{and} \quad j_1(x', s, \vartheta) := (x', s, \vartheta).$$

It follows from Lemma 6.3 that we can assume the primitive of the symplectic structure on Σ over $\nu_\varepsilon \Sigma_i'$ to be given by $\alpha_i = \alpha|_{\Sigma_i'} + s^2 d\vartheta$. Endow $\Sigma' \times (-1,1) \times S^1$ with an exact symplectic form whose primitive is given by $\beta = \alpha|_{\Sigma_i'} + f(s) d\vartheta$, where f is an appropriate function as in the proof of Theorem 6.1 (see [6, proof of Theorem 7.4.3], and note that we have $j_i^* \beta = \alpha_i$. Hence, the quotient carries a well-defined symplectic structure which coincides with the original one outside $\nu\Sigma_i'$.

Given the above description of the page, we can write the twist map $\mathcal{D}$ as a self-diffeomorphism of $\Sigma' \times (-1,1) \times S^1$ defined by

$$\mathcal{D}(x', s, \vartheta) := \big(x', s, \vartheta + \tau(\boldsymbol{r}, s) \big),$$

and extend it trivially to $\widetilde{\Sigma}$. Here, $\boldsymbol{r} = \boldsymbol{r}(x')$ and $\tau(\boldsymbol{r}, s) = -4\pi \sigma_1(\boldsymbol{r}) \sigma_2(\frac{s+1}{2})$, where σ_1 is a cut-off function interpolating between $\sigma_1(0) = 0$ and 1, and σ_2 is a bump function with $\sigma_2(0) = 1$ and vanishing near -1 and 1. The twist map $\mathcal{D}$ is not symplectic as one computes

$$\mathcal{D}^* d\beta = d\beta + d(f(s) d\tau).$$

However, this is equal to $d\alpha + f' ds \wedge d\vartheta - 4\pi f'(s) \sigma_2(\frac{s+1}{2}) \sigma_1'(r) ds \wedge dr$, so we have $(\mathcal{D}^* d\beta)^n = (d\beta)^n$. Similarly, for $\omega_t := d\beta + t \cdot d(f(s) d\tau)$ we have $\omega_t^n = (d\beta)^n$ and $[\omega_t] = 0$ for all $t \in \mathbb{R}$, i.e. ω_t is a family of symplectic forms in the same cohomology class connecting $\mathcal{D}^* d\beta$ and $d\beta$. Since the deformation is compactly supported, we can apply a Moser stability theorem for symplectic structures (cf. [13, Theorem 3.17]) to obtain an isotopy φ_t with $\varphi_t^* \omega_t = \omega_0$. Hence, $\varphi_1 \circ \mathcal{D}$ is symplectic and smoothly isotopic to $\mathcal{D}$, i.e. $\Big(\widetilde{\Sigma},\, \phi \circ \psi \circ (\varphi_1 \circ \mathcal{D}) \Big)$ is an abstract open book description of the binding sum $\#_B M$, where $\phi \circ \psi$ is the monodromy of the abstract open book realising the push-offs as nested abstract open book described in Proposition 6.4. Furthermore, the abstract open book is endowed with a contact structure which coincides with the original contact structure away from the area of identification.

It is worth noting that the handle interpretation of the resulting page of the open book decomposition of the binding sum can also be adapted to the contact case, i.e. the new page can be obtained from the original page by the consecutive attachment of two generalised *symplectic* 1-handles as introduced in [11].

References

[1] J. W. Alexander, A lemma on systems of knotted curves, *Proc. Nat. Acad. Sci. U.S.A.* **9** (1923), 93–95.

[2] T. Bröcker and K. Jänich, Einführung in die Differentialtopologie, *Springer-Verlag*, Berlin (1973).

[3] M. Dörner, The Space of Contact Forms Adapted to an Open Book, *Ph.D. thesis*, University of Cologne (2014).

[4] Y. Eliashberg, Topological characterization of Stein manifolds of dimension > 2, *Internat. J. Math.* **1** (1990), 29–46.

[5] J. Etnyre, Lectures on open book decompositions and contact structures, `arXiv:math/0409402`.

[6] H. Geiges, An Introduction to Contact Topology, *Cambridge University Press*, Cambridge (2008).

[7] E. Giroux, Géométrie de contact: de la dimension trois vers les dimensions supérieures, *Proceedings of the International Congress of Mathematicians Vol. II (Beijing, 2002)*, Higher Education Press, Beijing (2002), 405–414.

[8] E. Giroux, What is... an open book?, *Notices Amer. Math. Soc.* **52** (2005).

[9] E. Giroux and J.-P. Mohsen, Structures de contact et fibrations symplectiques au-dessus du cercle, *in preparation*.

[10] M. Klukas, Open book decompositions of fibre sums in contact topology, *Algebr. Geom. Topol.* **16** (2016), 1253–1277.

[11] M. Klukas, Open books and exact symplectic cobordisms, `arXiv:1207.5647`.

[12] T. Lawson, Open book decompositions for odd dimensional manifolds, *Topology* **17** (1978), 189–192.

[13] D. McDuff and D. Salamon, Introduction to Symplectic Topology, *The Clarendon Press*, New York (1998).

[14] F. Quinn, Open book decompositions, and the bordism of automorphisms, *Topology* **18** (1979), 55–73.

[15] A. Ranicki, High-Dimensional Knot Theory – Algebraic Surgery in Codimension 2, with an appendix by E. Winkelnkemper, *Springer Monogr. Math.*, Springer-Verlag, Berlin (1998).

[16] O. van Koert, Lecture notes on stabilization of contact open books, `arXiv:1012.4359`.

[17] A. Weinstein, Contact surgery and symplectic handlebodies, *Hokkaido Math. J.* **20** (1991), 241–251.

[18] H. E. Winkelnkemper, Manifolds as open books, *Bull. Amer. Math. Soc.* **79** (1973), 45–51.

(S. Durst) Mathematisches Institut, Universität zu Köln, Weyertal 86–90, 50931 Köln, Germany

E-mail address: `sdurst@math.uni-koeln.de`

(M. Klukas) Institute of Science and Technology Austria, Am Campus 1, A-3400 Klosterneuburg, Austria

E-mail address: `mirko.klukas@gmail.com`